

Klinefelter Syndrome—A Clinical Update

Kristian A. Groth, Anne Skakkebak, Christian Høst, Claus Højbjerg Gravholt, and Anders Bojesen

Departments of Molecular Medicine (K.A.G., C.H.G.) and Endocrinology and Internal Medicine (A.S., C.H., C.H.G.), Aarhus University Hospital, DK-8000 Aarhus C, Denmark; and Department of Clinical Genetics (A.B.), Vejle Hospital, Sygehus Lillebaelt, 7100 Vejle, Denmark

Context: Recently, new clinically important information regarding Klinefelter syndrome (KS) has been published. We review aspects of epidemiology, endocrinology, metabolism, body composition, and neuropsychology with reference to recent genetic discoveries.

Evidence Acquisition: PubMed was searched for “Klinefelter,” “Klinefelter’s,” and “XXY” in titles and abstracts. Relevant papers were obtained and reviewed, as well as other articles selected by the authors.

Evidence Synthesis: KS is the most common sex chromosome disorder in males, affecting one in 660 men. The genetic background is the extra X-chromosome, which may be inherited from either parent. Most genes from the extra X undergo inactivation, but some escape and serve as the putative genetic cause of the syndrome. KS is severely underdiagnosed or is diagnosed late in life, roughly 25% are diagnosed, and the mean age of diagnosis is in the mid-30s. KS is associated with an increased morbidity resulting in loss of approximately 2 yr in life span with an increased mortality from many different diseases. The key findings in KS are small testes, hypergonadotropic hypogonadism, and cognitive impairment. The hypogonadism may lead to changes in body composition and a risk of developing metabolic syndrome and type 2 diabetes. The cognitive impairment is mainly in the area of language processing. Boys with KS are often in need of speech therapy, and many suffer from learning disability and may benefit from special education. Medical treatment is mainly testosterone replacement therapy to alleviate acute and long-term consequences of hypogonadism as well as treating or preventing the frequent comorbidity.

Conclusions: More emphasis should be placed on increasing the rate of diagnosis and generating evidence for timing and dose of testosterone replacement. Treatment of KS should be a multidisciplinary task including pediatricians, speech therapists, general practitioners, psychologists, infertility specialists, urologists, and endocrinologists. (*J Clin Endocrinol Metab* 98: 20–30, 2013)

Klinefelter syndrome, 47,XXY (KS), occurs in about 150 per 100,000 males and is the most frequent chromosomal aberration in males. It was first described in 1942 (1), with a number of additional conditions, characteristics, and abnormalities described in later publications. KS has a genetic background, with characteristics involving numerous specialties such as embryology, pediatrics, endocrinology, cardiology, psychology, psychiatry, urology, and epidemiology. We have lately expanded our knowledge concerning KS.

This review is on aspects of epidemiology, endocrinology, metabolism, cardiology, body composition, and neuropsychology of the syndrome, with reference to recent genetic discoveries.

Diagnosis, Epidemiology, and Genetics

The designation “Klinefelter syndrome” is a clinical characterization. No firm guidelines for the diagnosis exist, but most agree that the cardinal stigmata include small testes

in virtually all KS, hypergonadotropic hypogonadism, gynecomastia, learning difficulties, and infertility. However, the clinical presentation of XXY males may appear in many cases to be similar to that of XY males, and thus it is difficult to make a diagnosis of KS without karyotyping. A number of congenital malformations and conditions are often seen in KS (Table 1).

The genetic background for the KS phenotype is based on the presence of the extra X-chromosome. The genetic disorder is also seen in domestic and wild animals (2). As in women, one of the extra X-chromosomes undergoes inactivation, and the phenotype is presumed to be the consequence of the presence of the non-inactivated extra genes from the X-chromosome, although other genetic mechanisms are possible. Of these genes, the only one that has been clearly shown to influence the phenotype in KS is

the short-stature homeobox-containing gene on chromosome X (*SHOX*) situated in the pseudoautosomal region 1 on Xp. *SHOX* haploinsufficiency has been implicated in growth retardation and bone changes in Turner syndrome and Leri-Weill dyschondrosteosis (3, 4) and is also implicated in the slightly accelerated growth in Klinefelter, 47,XXX and 47,XXY syndrome (5). Brain natriuretic peptide and fibroblast growth factor receptor 3 are transcriptional targets of *SHOX* (6, 7), and this knowledge may enhance our understanding of the phenotypic consequences of the syndrome. The CAG repeat number in the androgen receptor seems to be related to some phenotypic traits in KS, like height and hematocrit and possibly others (8–10), whereas it is more doubtful whether parental origin of the extra X-chromosome influences phenotype. More than 10% of the genes located on the X-chromosome are expressed in the testis and therefore likely to play a role in KS (11).

In the first publication, the syndrome was described as “not uncommon” (1), but it was not until large-scale chromosome analyses in newborns were performed that the “true” prevalence was established, however with great variation. By pooling the data from eight different studies, we estimated the prevalence to be 152 per 100,000 live-born males, confirmed by the prevalence in prenatally diagnosed KS in Denmark (12) and a large screening study from Georgia where DNA from dried blood spots from 36,124 newborn boys showed a prevalence of 158 per 100,000 (13). An increase in the prevalence has been proposed (14), and the prevalence of KS may differ between populations, exemplified by a recent Australian study finding a prevalence of 223 per 100,000 (15).

Establishing the prevalence has enabled us to study the incidence of diagnosis. In Denmark, only 25% of the expected numbers are diagnosed, and barely 10% of these are diagnosed before puberty, indicating a severe delay in diagnosis, underdiagnosis, and nondiagnosis (12). Similar frequency of diagnosis was estimated from a British study finding approximately 100 of estimated 525 each year (16), but in Australia the diagnostic activity seems greater because they find approximately 50% of the expected numbers (15). The reason so many KS men will go through their lives without a diagnosis is unknown, but the relatively mild phenotype is a tempting explanation. The hidden proportion of nondiagnosed patients should be kept in mind when looking at data concerning KS because ascertainment bias is always present, apart from screening studies in the newborn.

Epidemiological studies in KS have been conducted in two major cohorts, the British and the Danish. Data concerning mortality and cancer incidence have been published from the British cohort (16, 17). In the Danish co-

TABLE 1. Abnormalities associated with KS and their tentative frequencies^a

Feature	Frequency (%)
Infertility (adults) (8, 57)	91–99 ^b
Small testes (bi-testicular size <6 ml) (8)	>95
Increased gonadotropin levels (57)	>95
Azoospermia (adults) (57)	>95
Learning disabilities (children) (74)	>75
Decreased testosterone levels (57)	63–85
Decreased facial hair (adults) (57)	60–80
Decreased pubic hair (adults) (57)	30–60
Gynecomastia (adolescents, adults) (8, 33, 74)	38–75
Delay of speech development (children) (74)	40
Increased height (prepubertal, adults) (74, 123)	30
Abdominal adiposity (adults) (36)	~50
Metabolic syndrome (adults) (36)	46
Osteopenia (adults) (51, 124)	5–40
Type 2 diabetes (adults) (19, 36)	10–39
Cryptorchidism (8, 74)	27–37
Decreased penile size (children) (74)	10–25
Psychiatric disturbances (children) (74)	25
Congenital malformations, cleft palate, inguinal hernia (125)	~18
Osteoporosis (adults) (124)	10
Mitral valve prolapse (adults) (126, 127)	0–55
Breast cancer (adults) (16, 128)	Increased risk (~50 fold)
Mediastinal cancers (children) (22)	increased risk (~500 fold)
Fractures (17, 18)	Increased risk (2–40 fold)

^a The hidden proportion of nondiagnosed patients should be kept in mind when studying the estimates given in this table. Ascertainment bias is present in all studies presented so far, and no separate study has presented more than 25% of a given population of KS individuals.

^b Lanfranco *et al.* (8) found 8.4% of males with sperm in the ejaculate. Whether this translates into 8.4% of KS males being able to achieve a normal pregnancy is doubtful.

hort, mortality, morbidity, socioeconomic data, and criminality have been studied (18–21). Together, these studies show clearly that the average KS individual who comes to clinical diagnosis fares less well than his peers. The expected life span was found to be reduced by 1.5 to 2 yr, with increased mortality from a range of different disorders including diabetes, lung diseases, epilepsy, cerebrovascular disease, and vascular insufficiency of the intestine (17, 18). Mortality from breast cancer was greatly increased in the British cohort, as well as mortality from non-Hodgkin lymphoma and lung cancer (16). In a former study in the Danish cohort, the risk of breast cancer was not increased, but the risk of mediastinal tumor was greatly increased (22). The risk of being admitted to hospital was increased by 70% overall, and the highest risks of being admitted were due to congenital malformations and psychiatric, endocrine, and metabolic disorders (19). In seeking to understand the basis for the increased morbidity and mortality, we studied the socioeconomic profile of the KS population compared with a large background population and found that KS men are characterized by shorter education, lower income, earlier retirement, more unemployment, and entering marriage less frequently and with fewer offspring. Mortality among KS men was significantly increased (hazard ratio, 1.9) and remained so after adjustment for cohabitation and educational status (hazard ratio, 1.5), indicating that socioeconomic parameters may explain some but not all the excess mortality in KS (20).

During the 1960s and 1970s, increased prevalence of KS and 47,XYY was found in prisons and institutions for mentally handicapped individuals, and a general increased rate of criminal behavior and an increased crime rate were reported among both cohorts, especially due to sexual crimes (23, 24). These studies were criticized because of ascertainment bias because they investigated institutionalized individuals and were hence disregarded as prejudicial. We wanted to investigate crime rates in our cohort of nearly 1000 KS men compared with a control group of nearly 100,000 age-matched men, simply to describe whether the prejudice of increased criminality was true or false. To our surprise, it was true; even after adjustment for some socioeconomic parameters, the crime rate for sexual abuse and arson was still significantly increased, whereas traffic offenses and drug-related crime were significantly decreased (21).

The Pituitary Testicular Axis

Males with KS are considered infertile; however, recent studies have shown that modern technology such as tes-

ticular sperm extraction followed by intracytoplasmic sperm injection can allow fatherhood in KS (25), and many couples also opt for adoption or the use of donor semen as a means of becoming parents (20). The most recent results with advanced technology show a sperm recovery rate of 66% (which is euploid in the vast majority), and 45% of these achieved a live birth of a child (26). If an increase in circulating testosterone, and thus also testicular testosterone, was achieved by treatment with either human chorionic gonadotropin or aromatase inhibitors, better results were observed (26). There are also indications that sperm may decline or disappear with age (27), which could be an argument for early retrieval of sperm by modern techniques. However, the pathophysiological background for the development of infertility and hypogonadism is still poorly understood. The “minipuberty” seen in infants during the first 3 months of life, with a surge in testosterone, was at one point described as blunted in KS infants (28, 29); however, a more recent study did not confirm this finding (30). Typical testicular histology in KS is with hyalinization of seminiferous tubules with loss of germ cells and Leydig cell hyperplasia; although focal spermatogenesis may be found with the possibility of surgical extraction of viable sperm (25). The hyalinization of the seminiferous tubules probably occurs at midpuberty (31), which is usually timed correctly (32) with bi-testicular growth to approximately 6 ml and shrinkage thereafter to a pathological adult size of less than 6 ml as examined by ultrasonography (8). At the beginning of puberty, the levels of FSH, LH, and testosterone are normal, but FSH and LH start to increase and testosterone to decline compared with normal boys (33). In adult KS patients, levels of testosterone, insulin-like factor 3 (34), inhibin B (32), and anti-Müllerian hormone (35) are decreased, whereas FSH and LH are elevated and 17 β -estradiol and SHBG are comparable to controls. The ensuing hypogonadism is relative rather than absolute, with the typical level of testosterone in the low normal range or subnormal (8, 36). Sexual dysfunction in KS is probably high, but it seems to be linked to low testosterone (37).

Glucose Homeostasis and Physical Fitness

An association between KS and diabetes has long been recognized. In 1969, Nielsen *et al.* (38) described an increased prevalence (39%) of a diabetic oral glucose tolerance test, whereas others found decreased insulin sensitivity and elevated fasting insulin levels in seven patients with KS (39). We recently described a strikingly high incidence of the metabolic syndrome and insulin resistance

in 70 patients with KS compared with an age-matched control group. Almost half of the KS patients fulfilled criteria for the metabolic syndrome, whereas it was true for only 10% of the control group (36), findings corroborated by Ishikawa *et al.* (40). Most recently, a study on 89 prepubertal KS boys found 37% to have elevated low-density lipoprotein cholesterol, 24% with insulin resistance, and 7% meeting the criteria of the metabolic syndrome (41).

Cross-sectional studies have consistently reported an inverse relationship between plasma testosterone and insulin resistance in normal males (42). Type 2 diabetes is frequent in hypogonadal patients (19), and vice versa, hypogonadism is also more prevalent among type 2 diabetics with presumed normal karyotype than in age-matched controls (42). These findings are minimized (43), or even absent, in some studies (36, 44) when correcting for body mass index (BMI) or waist to hip ratio, and question whether this association is largely mediated by adiposity rather than testosterone itself. Indeed, testosterone treatment of hypogonadal patients with type 2 diabetes primarily improves insulin sensitivity in obese patients (45), but not in lean patients (46). This indicates that improvements in insulin sensitivity after therapy largely depend on the amount of “modifiable fat,” especially visceral fat. Conversely, in a recent meta-analysis encompassing several cross-sectional studies and some 2900 men, of whom 850 had type 2 diabetes, testosterone levels were significantly lower in type 2 diabetics even after controlling for age, BMI, and waist to hip ratio, whereas high testosterone levels were associated with a decreased risk of type 2 diabetes mellitus (47).

Indeed, there are indications that testosterone itself has direct effects on insulin sensitivity. In patients with hypogonadotropic hypogonadism, cessation of testosterone replacement therapy resulted in deteriorated insulin sensitivity within only 14 d (48). Other studies using a hyperinsulinemic euglycemic clamp found no effect on insulin sensitivity during short-term hypogonadism in healthy controls (49), whereas 1-wk treatment of healthy lean men with aromatase inhibitors resulted in slight supraphysiological testosterone levels and improved insulin sensitivity (50).

KS patients have altered body composition with increased total body and truncal fat (36) and decreased lean body mass accompanied by lower aerobic capacity and muscle strength in both biceps and quadriceps muscles (51). At present, no studies have examined the effects of testosterone treatment on muscle strength or other measures of physical fitness in KS patients. In a recent cross-sectional study of elderly men, low testosterone by multivariate analysis was associated with lower grip strength and hemoglobin, but not with 4-m walking speed or muscle mass (52). In hypogonadal elderly men, testosterone

has been shown to increase hand-grip strength and physical performance during 36-month treatment (53) and to improve lower and upper body muscle strength (54). In addition, results from a recent randomized, placebo-controlled study suggest that testosterone treatment may prevent the age-related loss of muscle mass, strength, and physical function, and that it may improve quality of life in frail elderly men with low to borderline-low testosterone (55).

Combined, both epidemiological and clinical studies show clear evidence of a dramatically increased risk of diabetes and metabolic syndrome in KS. Available evidence does not support testosterone replacement therapy of KS patients with the aim of improving insulin sensitivity, but such effects may occur indirectly through favorable effects on body composition and physical fitness, although formal studies are not at hand in these patients.

Anthropometry and Body Composition

The KS phenotype varies greatly, and there is no exclusive symptom to define the syndrome. This might be one of the reasons why the syndrome is highly underdiagnosed, with less than 25% of adult male patients diagnosed (12). Many different phenotypic abnormalities have been associated with KS (Table 1). KS patients have an accelerated growth from early childhood and tend to become taller than controls. Final height tends to be greater in patients who initiated testosterone therapy after the age of 18 compared with patients treated from the time of puberty (56). The increased height is mainly attributed to abnormally long legs (8, 57). Although KS patients are abdominally obese, BMI may often be in the normal range due to an unfavorable muscle/fat ratio with decreased muscle mass and increased body fat, along with greatly elevated leptin levels (36). Greater body fat mass, however, is already present before puberty, pointing toward genetic influences on body fat in KS (58). In addition, we know that androgens can prevent the differentiation of pluripotent cells into an adipogenic lineage (59), whereas hypogonadism independently predicts development of abdominal adiposity in men with normal chromosomes (60). Conversely, testosterone treatment causes dose-dependent changes in fat-free mass (which inversely relates to increased testosterone levels), and treatment of middle-aged abdominally obese men reduces the amount of intraabdominal fat (61). Recently, a study in KS patients showed that testosterone treatment only partially corrected the unfavorable muscle/fat ratio, but these findings may be a result of the insufficient testosterone doses used (58). Lower levels of testosterone, free testosterone, and SHBG are found in obese

men compared with nonobese men (62), but the mechanism for lower testosterone in obese men remains unclear. In fact, it is still unclear how hypogonadism leads to abdominal adiposity and how the abdominal adiposity leads to decreased testosterone production, but both scenarios may be part of a self-perpetuating vicious circle—the so-called hypogonadal-obesity circle (63).

Despite the close relationship between testosterone and body composition, it is uncertain whether hypogonadal males and KS patients are comparable or whether genetic factors related to KS add another layer of complexity. Prospective studies in KS are needed to address these issues and perhaps clarify whether early androgen deficiency predisposes to the hypogonadal-obese insulin-resistant phenotype of KS, or whether other factors related to the sex chromosome trisomy are responsible. Such information will help clinicians decide the optimal timing and mode of hormone therapy to these patients.

Bone mineralization

Decreased bone mass (51, 64–66) and outright fractures and osteoporosis affecting morbidity but also mortality (17–19) have long been linked to KS. This propensity toward low bone mass is readily explained by hypogonadism, fuelling low physical exercise capacity and muscle strength (Fig. 1), and treatment with testosterone improves bone mineral density (BMD) (65), but future studies are needed to provide evidence for a positive effect on fractures and the development of osteoporosis as well. Speculatively, changes in thyroid functioning may exacerbate low BMD (67, 68).




FIG. 1. Significant correlation between muscle strength in the arm (biceps) and BMD in the distal forearm in both KS (●, solid line, $r^2 = 0.19$; $P = 0.001$) and healthy subjects (○, dashed line, $r^2 = 0.17$; $P < 0.001$). [Reprinted from A. Bojesen et al.: Bone mineral density in Klinefelter syndrome is reduced and primarily determined by muscle strength and resorptive markers, but not directly by testosterone. *Osteoporos Int* 22:1441–1450, 2011 (51), with permission. © Springer.]

Neurocognitive function

The neuropsychological phenotype in boys and adults with KS is highly variable. The most radically impaired area is the verbal, which includes delay in early language development (69–75), learning disabilities in reading and spelling (74, 76–78), difficulties in production of syntax (78), word retrieval (78, 79), and nonsemantic cues in spoken language (80), resulting in a high degree of these boys getting speech and language therapy and special help in school (74, 79, 81–83). These disabilities in verbal cognitive function are probably part of the explanation for adults with KS being found to have a lower educational level (20, 84). The general cognitive ability seems to be close to normal levels, with a mean IQ of 87.9–110 (78, 85–94). Other cognitive problems include impairments in both verbal (86, 87) and nonverbal memory (78, 95) and in executive functions (90, 91, 95–98). Visual and spatial cognitive abilities are normal in boys and adults with KS (79, 87–89, 91), and arithmetic abilities are normal (89) or mildly impaired (83, 87, 97, 99).

A positive effect of testosterone therapy has been seen on behavior (100, 101), energy level (100), well-being (100), learning capacity (100, 101), and verbal fluency in patients with KS (102), whereas two other studies did not report any effects of testosterone treatment (89, 103). Data concerning the effect of testosterone therapy on brain volumes are sparse. A significant reduction in left temporal lobe gray matter volumes in untreated KS patients has been reported in one study (102), but this was not supported by another study (89). Thus, more studies are needed to investigate the effect of testosterone therapy on cognition and brain morphology. Several genetic mechanisms have been proposed to contribute to the neuropsychological phenotype, which include skewed X-chromosome inactivation, parental origin of the supernumerary X-chromosome, polymorphism of the androgen receptor gene CAG repeat, and a gene dosage effect based on the presence of the extra X-chromosome. Skewed X-chromosome inactivation occurs in 9–21% of KS patients; however, due to the small number of KS subjects with skewed X-chromosome, no information on this is available (103). Regarding parental origin of the supernumerary X-chromosome, one study reported significantly higher speech and language problems in the group with paternal origin (104); however, other studies do not support these findings (70, 103). Short androgen receptor CAG repeat length was associated with stable partnership and professions that require higher educational levels (9); however, an association between the androgen receptor gene polymorphism and neuropsychological outcome is not supported by other studies (103, 104). A recent study found differentially expressed genes in KS patients showing significant correlations with verbal cognitive test scores (105). These interesting results need further confirmation by other studies.

TABLE 2. Testosterone preparations available and suggested dosages for adults

Substance	Brand name (manufacturer)	Format	Route of administration	Suggested dose
Testosterone undecanoate	Andriol (Organon, Oss, The Netherlands)	40-mg capsule	Oral	120–160 mg/d
Testosterone undecanoate	Nebido (Schering, Berlin, Germany)	1000-mg injection	Intramuscular	1000 mg every 9–16 wk
Testosterone enanthate	Testoviron (Schering, Berlin, Germany)	250-mg injection	Intramuscular	250 mg every 2–4 wk
Testosterone	Testim (Ipsen, Paris, France)	Gel	Skin	50 mg/d
Testosterone	Testogel (Laboratoires Besins, Paris, France)	Gel	Skin	50 mg/d
Testosterone	Tostran (ProStrakan, Galashiels, UK)	Gel	Skin	40–60 mg/d
Testosterone	Implants (Organon, Oss, The Netherlands)	Pellets	Subcutaneous	400–800 mg every 4–6 months
Testosterone	Striant (Columbia Laboratories, Livingston, NJ)	Buccal adhesive	Buccal	60 mg/d
Testosterone	Androderm (Watson Pharma Inc., Corona, CA)	Transdermal patch	Skin	5–15 mg/d
Testosterone	Testoderm (Alza Corp., Mountain View, CA)	Transdermal patch	Scrotal skin	2.5–10.0 mg/d

For children and adolescents, lower doses should be given (118). Some preparations are not available in all countries.

Psychiatric morbidity

A review of studies of male psychiatric inpatients from the 1960s to the 1990s found a frequency of KS ranging from 0 to 4.8% (mean, 0.8%) among schizophrenic patients (106), a 4- to 5-fold increase compared with the prevalence of KS in the general population. Prospective studies of KS also reported higher rates of psychiatric referral among boys with KS (74), and in adolescence, 54% of KS males had mild to moderate psychiatric disorders (77). A register study reported that individuals with KS have an increased hazard ratio of 3.65 (95% confidence interval, 2.92–4.55) of being hospitalized with a psychotic disorder (19). Studies on unselected KS boys and adults for psychiatric disorders support this by finding a significantly increased prevalence of schizotypal traits, schizophrenia symptoms, psychotic disorders (93, 95, 107, 108), depressive disorders (107, 108), anxiety disorder (108), autism spectrum diseases (94, 108, 109), and attention deficit/hyperactivity disorders (91, 92, 108). Recently, two studies have investigated the relationship between parent-of-origin of the extra X-chromosome and the psychopathology seen in KS patients (107, 108) and found conflicting results.

Clinical care

We believe that treatment and care of patients with KS is a multidisciplinary task that ideally should involve speech therapists, psychologists, general practitioners, pediatricians, endocrinologists, urologists, and infertility specialists. Infants with KS are rarely diagnosed because they lack KS-specific stigmata. Rarely, however, KS boys have micropenis, which can be treated successfully with topical testosterone cream or single injections with im testosterone (110). The most serious problem in early childhood is the delay of speech development and learning

problems affecting perhaps half of the boys with KS (74). Careful observation is needed to refer these boys to speech therapists if delay of speech is observed. The same holds true for learning disabilities that were observed in 77% of boys with KS followed from birth to adulthood (74), and it is necessary to develop schemes to enhance learning in schools.

TABLE 3. Outpatient program for patients with KS

At baseline
Confirmation of karyotype, if necessary
Sex hormones: testosterone, estrogen, SHBG, FSH, and LH
Fasting glucose, lipids, and HbA1c
Thyroid status, hemoglobin, hematocrit
Physical examination including BP, height, weight, waist, testes, gynecomastia, and varicose veins
Bone densitometry (DEXA scan) and vitamin D status, p-calcium
Information about the syndrome
Initiation of androgen treatment (injections, transdermal, or oral)
Questions about well-being, physical activity, energy, sexual activity, libido, socioeconomic situation
Echocardiography if deemed necessary
Discussion of fertility issues often resulting in referral to a fertility clinic
Consider referral to plastic surgeon for correction of gynecomastia
Consider referral to psychologist
Annual (every 3 months initially)
Physical examination including BP, height, weight, waist, and gynecomastia
Sex hormones: testosterone, estrogen, SHBG, FSH, and LH (nadir values)
Fasting glucose, lipids, and HbA1c
Thyroid status, hemoglobin, hematocrit
Questions about well-being, physical activity, energy, sexual activity, libido
Every second year or up to every tenth year
Bone densitometry (DEXA scan) and vitamin D status, p-calcium

HbA1c, Glycosylated hemoglobin; BP, blood pressure; DEXA, dual-energy x-ray absorptiometry.




FIG. 2. The effect of increased dosage of genes on the X-chromosome (and possibly other genetic mechanisms, such as a different methylation pattern and skewed X-inactivation working in concert) is at the center of the current understanding of pathogenesis in KS. This leads to a range of effects affecting the endocrine system, with especially hypergonadotropic hypogonadism, changed body composition, and increased risk of osteoporosis. Hypogonadism has pervasive effects, affecting: 1) different hormone levels; 2) cardiovascular features; 3) metabolic features; and 4) features related to sex hormones, such as infertility. In addition, mounting evidence suggests that hypogonadism in KS leads to a vicious circle increasing insulin resistance and enhancing unfavorable body composition changes either directly or indirectly. Increasingly, awareness is growing concerning a poor socioeconomic profile and psychiatric morbidity. Arrows indicate possible consequences—not all interactions have been shown in scientific studies. BNP, Brain natriuretic peptide; FGFR3, fibroblast growth factor receptor 3; ADHD, attention deficit/hyperactivity disorder.

We suggest initiation of testosterone treatment at the beginning of puberty, as FSH and LH start to rise, to secure a proper masculine development of sexual characteristics but also to secure a sufficient increase in BMD and muscle bulk to prevent subsequent osteoporosis. It is very important to

TABLE 4. Major issues in KS

Problem	Possible solution
Will early diagnosis lead to better outcome?	Prospective screening studies with health technology assessment with reference to medical ethics
Late diagnosis and nondiagnosis	Examination of dried blood spots with new molecular genetic techniques
Poor learning in school	Early diagnosis leading to better learning schemes and perhaps early treatment with T
Effect of T	Randomized clinical trials with T and placebo with study of numerous variables
Type 2 diabetes	Randomized clinical trials with T and placebo
Poor socioeconomic outcome	Improvements in schooling and possibly early treatment
Increased morbidity	Improvements in adult care with multidisciplinary approach
Infertility	Improved understanding of pathophysiology of germ cell loss through animal models; better testicular sperm extraction techniques

T, Testosterone.

discuss fertility, and postponement of treatment may be an option if one wants to retrieve viable sperm at this stage. Testosterone treatment in pubertal KS boys has also been reported to increase energy and endurance and to improve mood, concentration, and relations to others (111), and it seems that there are increased psychosocial problems in periods without testosterone treatment in pubertal KS (112).

We advise lifelong treatment in order to prevent osteoporosis, obesity, metabolic syndrome, and diabetes. However, this practice is not evidence-based. Treatment in young hypogonadal men has been shown to have a positive impact also on fat mass, muscle mass, and muscle strength, as well as sexual activity and related areas, and it improves positive aspects of mood (113). In older hypogonadal males, there are also limited data to suggest positive effects of treatment on visuospatial cognition and verbal memory (114). Although some KS patients have normal testosterone values, virtually all have increased gonadotropin levels.

Some with KS may not realize that they have symptoms, and only after a trial period of treatment do they see the benefits of treatment. We believe that all KS patients should receive testosterone treatment if their gonadotropins are elevated, although their testosterone levels may be within the lower end of the normal range (115), using bivariate charts of testosterone vs. LH for proper dosage (116). Certainly, patients should be treated if they are suffering from hypogonadal symptoms (lack of energy, decreased libido, and also including abdominal adiposity, etc.). Treatment options include oral, transdermal, im, and buccal routes of administration (Table 2) (117). Transdermal gel preparations offer the best pharmacodynamic profile, but im injections remain popular due to ease of administration. Treatment of children and adolescents presents special problems, dose escalation must be considered (118), and we would usually start with oral or transdermal treatment. The aim of treatment should include normalization in LH and testosterone levels in the mid-normal range, rather than low-normal nadir values of testosterone, because our experience with KS patients is that most of the patients are insufficiently treated. It is not always feasible to normalize LH due to elevated hematocrit, and one should also focus on the subjective symptoms reported by the patient, especially to avoid high levels of testosterone that can occur with injection therapy and may cause discomfort.

Our clinical outpatient program for patients with KS is presented in Table 3. Initially, testosterone treatment should be followed with visits every 3 months until testosterone dose is adjusted, and thereafter annually.

Perspectives

Our perception of KS is changing, and the way we see KS today is multifaceted with much new knowledge being added in recent years (Fig. 2). A number of issues need to be resolved in KS (Table 4), and large, possibly international, collaborations are suggested in this regard. Not a single randomized, placebo-controlled study addressing the effects of testosterone in KS patients has been published. We propose randomized, placebo-controlled studies on adults with KS, with testosterone preparations that will restore testosterone to normal values, in a population large enough to detect small changes in BMD, body composition, insulin sensitivity, and also modalities of quality of life. The recently established animal models are exciting and could shed new light on fertility (119), bone morphology (120), and brain functioning and learning difficulties (121), as well as other issues. The delay in diagnosis and outright nondiagnosis is also problematic—we need to devise programs to improve the advent of early diagnosis, and we think that new approaches, such as pervasive testing of all neonatal dried blood spot samples with new molecular genetic techniques, should reduce the costs substantially in comparison with karyotyping (122). However, before such an approach is implemented, it is important to discuss whether or not early diagnosis will in fact lead to a better outcome, and to that end we dearly need more scientific data. We need large, prospective, nationwide screening studies with health technology assessment. Hopefully, future studies will provide the evidence that is essential for optimizing the treatment of KS patients.

Acknowledgments

Address all correspondence and requests for reprints to: Claus Højbjerg Gravholt, M.D., Ph.D., Department of Endocrinology and Internal Medicine, Aarhus University Hospital, DK-8000 Aarhus C, Denmark. E-mail: ch.gravholt@dadnet.dk.

This work was supported by grants from the Danish Health Research Council (Aarhus University-Novo Nordisk Center for Research in Growth and Regeneration, Grant 9600822), Aarhus University, the Lundbeck Foundation, the Aase and Einar Danielsen Foundation, the A. P. Møller and wife Chastine Mc-Kinney Møllers Foundation, the Danish Diabetes Association, Central Denmark Region, and the Danish Ministry of Science, Technology, and Innovation.

Disclosure Summary: The authors have nothing to disclose.

References

1. Klinefelter HF, Reifenstein EC, Albright F 1942 Syndrome characterized by gynecomastia, aspermatogenesis without a-Leydigism, and increased excretion of follicle-stimulating hormone. *J Clin Endocrinol* 2:615–627
2. Hauffe HC, Giménez MD, Garagna S, Searle JB 2010 First wild XXY house mice. *Chromosome Res* 18:599–604
3. Rao E, Weiss B, Fukami M, Rump A, Niesler B, Mertz A, Muroya K, Binder G, Kirsch S, Winkelmann M, Nordsiek G, Heinrich U, Breuning MH, Ranke MB, Rosenthal A, Ogata T, Rappold GA 1997 Pseudoautosomal deletions encompassing a novel homeobox gene cause growth failure in idiopathic short stature and Turner syndrome. *Nat Genet* 16:54–63
4. Ellison JW, Wardak Z, Young MF, Gehron Robey P, Laig-Webster M, Chiong W 1997 PHOG, a candidate gene for involvement in the short stature of Turner syndrome. *Hum Mol Genet* 6:1341–1347
5. Ottesen AM, Aksglaede L, Garn I, Tartaglia N, Tassone F, Gravholt CH, Bojesen A, Sørensen K, Jørgensen N, Rajpert-De Meyts E, Gerdes T, Lind AM, Kjaergaard S, Juul A 2010 Increased number of sex chromosomes affects height in a nonlinear fashion: a study of 305 patients with sex chromosome aneuploidy. *Am J Med Genet A* 152A:1206–1212
6. Marchini A, Häcker B, Marttila T, Hesse V, Emons J, Weiss B, Karperien M, Rappold G 2007 BNP is a transcriptional target of the short stature homeobox gene SHOX. *Hum Mol Genet* 16:3081–3087
7. Decker E, Durand C, Bender S, Rödelsperger C, Glaser A, Hecht J, Schneider KU, Rappold G 2011 FGFR3 is a target of the homeobox transcription factor SHOX in limb development. *Hum Mol Genet* 20:1524–1535
8. Lanfranco F, Kamischke A, Zitzmann M, Nieschlag E 2004 Klinefelter's syndrome. *Lancet* 364:273–283
9. Zitzmann M, Depenbusch M, Gromoll J, Nieschlag E 2004 X-chromosome inactivation patterns and androgen receptor functionality influence phenotype and social characteristics as well as pharmacogenetics of testosterone therapy in Klinefelter patients. *J Clin Endocrinol Metab* 89:6208–6217
10. Bojesen A, Hertz JM, Gravholt CH 2011 Genotype and phenotype in Klinefelter syndrome—impact of androgen receptor polymorphism and skewed X inactivation. *Int J Androl* 34:e642–e648
11. Ross MT, Grafham DV, Coffey AJ, Scherer S, McLay K, Muzny D, Platzer M, Howell GR, Burrows C, Bird CP, Frankish A, Lovell FL, Howe KL, Ashurst JL, Fulton RS, Sudbrak R, Wen G, Jones MC, Hurler ME, Andrews TD, Scott CE, Searle S, Ramser J, Whittaker A, Deadman R, Carter NP, Hunt SE, Chen R, Cree A, Gunaratne P, Havlak P, Hodgson A, Metzker ML, Richards S, Scott G, Steffen D, Sodergren E, Wheeler DA, Worley KC, et al. 2005 The DNA sequence of the human X chromosome. *Nature* 434:325–337
12. Bojesen A, Juul S, Gravholt CH 2003 Prenatal and postnatal prevalence of Klinefelter syndrome: a national registry study. *J Clin Endocrinol Metab* 88:622–626
13. Coffee B, Keith K, Albizua I, Malone T, Mowrey J, Sherman SL, Warren ST 2009 Incidence of fragile X syndrome by newborn screening for methylated FMR1 DNA. *Am J Hum Genet* 85:503–514
14. Morris JK, Alberman E, Scott C, Jacobs P 2008 Is the prevalence of Klinefelter syndrome increasing? *Eur J Hum Genet* 16:163–170
15. Herlihy AS, Halliday JL, Cock ML, McLachlan RI 2011 The prevalence and diagnosis rates of Klinefelter syndrome: an Australian comparison. *Med J Aust* 194:24–28
16. Swerdlow AJ, Schoemaker MJ, Higgins CD, Wright AF, Jacobs PA 2005 Cancer incidence and mortality in men with Klinefelter syndrome: a cohort study. *J Natl Cancer Inst* 97:1204–1210
17. Swerdlow AJ, Higgins CD, Schoemaker MJ, Wright AF, Jacobs PA 2005 Mortality in patients with Klinefelter syndrome in Britain: a cohort study. *J Clin Endocrinol Metab* 90:6516–6522
18. Bojesen A, Juul S, Birkebaek N, Gravholt CH 2004 Increased mor-

- tality in Klinefelter syndrome. *J Clin Endocrinol Metab* 89:3830–3834
19. Bojesen A, Juul S, Birkebaek NH, Gravholt CH 2006 Morbidity in Klinefelter syndrome: a Danish register study based on hospital discharge diagnoses. *J Clin Endocrinol Metab* 91:1254–1260
 20. Bojesen A, Stochholm K, Juul S, Gravholt CH 2011 Socioeconomic trajectories affect mortality in Klinefelter syndrome. *J Clin Endocrinol Metab* 96:2098–2104
 21. Stochholm K, Bojesen A, Jensen AS, Juul S, Gravholt CH 2012 Criminality in men with Klinefelter's syndrome and XYY syndrome: a cohort study. *BMJ Open* 2:e000650
 22. Hasle H, Mellemegaard A, Nielsen J, Hansen J 1995 Cancer incidence in men with Klinefelter syndrome. *Br J Cancer* 71:416–420
 23. Schröder J, de la Chapelle A, Hakola P, Virkkunen M 1981 The frequency of XYY and XXY men among criminal offenders. *Acta Psychiatr Scand* 63:272–276
 24. Finley WH, McDanal Jr CE, Finley SC, Rosecrans CJ 1973 Prison survey for the XYY karyotype in tall inmates. *Behav Genet* 3:97–100
 25. Schiff JD, Palermo GD, Veeck LL, Goldstein M, Rosenwaks Z, Schlegel PN 2005 Success of testicular sperm injection and intracytoplasmic sperm injection in men with Klinefelter syndrome. *J Clin Endocrinol Metab* 90:6263–6267
 26. Ramasamy R, Ricci JA, Palermo GD, Gosden LV, Rosenwaks Z, Schlegel PN 2009 Successful fertility treatment for Klinefelter's syndrome. *J Urol* 182:1108–1113
 27. Wikström AM, Høi-Hansen CE, Dunkel L, Rajpert-De Meyts E 2007 Immunoreexpression of androgen receptor and nine markers of maturation in the testes of adolescent boys with Klinefelter syndrome: evidence for degeneration of germ cells at the onset of meiosis. *J Clin Endocrinol Metab* 92:714–719
 28. Lahlou N, Fennoy I, Carel JC, Roger M 2004 Inhibin B and anti-Müllerian hormone, but not testosterone levels, are normal in infants with nonmosaic Klinefelter syndrome. *J Clin Endocrinol Metab* 89:1864–1868
 29. Ross JL, Samango-Sprouse C, Lahlou N, Kowal K, Elder FF, Zinn A 2005 Early androgen deficiency in infants and young boys with 47,XXY Klinefelter syndrome. *Horm Res* 64:39–45
 30. Aksglaede L, Petersen JH, Main KM, Skakkebaek NE, Juul A 2007 High normal testosterone levels in infants with non-mosaic Klinefelter's syndrome. *Eur J Endocrinol* 157:345–350
 31. Wikström AM, Raivio T, Hadziselimovic F, Wikström S, Tuuri T, Dunkel L 2004 Klinefelter syndrome in adolescence: onset of puberty is associated with accelerated germ cell depletion. *J Clin Endocrinol Metab* 89:2263–2270
 32. Christiansen P, Andersson AM, Skakkebaek NE 2003 Longitudinal studies of inhibin B levels in boys and young adults with Klinefelter syndrome. *J Clin Endocrinol Metab* 88:888–891
 33. Salbenblatt JA, Bender BG, Puck MH, Robinson A, Faiman C, Winter JS 1985 Pituitary-gonadal function in Klinefelter syndrome before and during puberty. *Pediatr Res* 19:82–86
 34. Bay K, Hartung S, Ivell R, Schumacher M, Jørgensen D, Jørgensen N, Holm M, Skakkebaek NE, Andersson AM 2005 Insulin-like factor 3 serum levels in 135 normal men and 85 men with testicular disorders: relationship to the luteinizing hormone-testosterone axis. *J Clin Endocrinol Metab* 90:3410–3418
 35. Aksglaede L, Christiansen P, Sørensen K, Boas M, Linneberg A, Main KM, Andersson AM, Skakkebaek NE, Juul A 2011 Serum concentrations of anti-Müllerian hormone (AMH) in 95 patients with Klinefelter syndrome with or without cryptorchidism. *Acta Paediatr* 100:839–845
 36. Bojesen A, Kristensen K, Birkebaek NH, Fedder J, Mosekilde L, Bennett P, Laurberg P, Frystyk J, Flyvbjerg A, Christiansen JS, Gravholt CH 2006 The metabolic syndrome is frequent in Klinefelter's syndrome and is associated with abdominal obesity and hypogonadism. *Diabetes Care* 29:1591–1598
 37. Corona G, Petrone L, Paggi F, Lotti F, Boddi V, Fisher A, Vignozzi L, Balercia G, Sforza A, Forti G, Mannucci E, Maggi M 2010 Sexual dysfunction in subjects with Klinefelter's syndrome. *Int J Androl* 33:574–580
 38. Nielsen J, Johansen K, Yde H 1969 Frequency of diabetes mellitus in patients with Klinefelter's syndrome of different chromosome constitutions and the XYY syndrome. Plasma insulin and growth hormone level after a glucose load. *J Clin Endocrinol Metab* 29:1062–1073
 39. Pei D, Sheu WH, Jeng CY, Liao WK, Fuh MM 1998 Insulin resistance in patients with Klinefelter's syndrome and idiopathic gonadotropin deficiency. *J Formos Med Assoc* 97:534–540
 40. Ishikawa T, Yamaguchi K, Kondo Y, Takenaka A, Fujisawa M 2008 Metabolic syndrome in men with Klinefelter's syndrome. *Urology* 71:1109–1113
 41. Bardsley MZ, Falkner B, Kowal K, Ross JL 2011 Insulin resistance and metabolic syndrome in prepubertal boys with Klinefelter syndrome. *Acta Paediatr* 100:866–870
 42. Grossmann M, Thomas MC, Panagiotopoulos S, Sharpe K, Macisaac RJ, Clarke S, Zajac JD, Jerums G 2008 Low testosterone levels are common and associated with insulin resistance in men with diabetes. *J Clin Endocrinol Metab* 93:1834–1840
 43. Grossmann M, Gianatti EJ, Zajac JD 2010 Testosterone and type 2 diabetes. *Curr Opin Endocrinol Diabetes Obes* 17:247–256
 44. Selvin E, Feinleib M, Zhang L, Rohrmann S, Rifai N, Nelson WG, Dobs A, Basaria S, Golden SH, Platz EA 2007 Androgens and diabetes in men: results from the Third National Health and Nutrition Examination Survey (NHANES III). *Diabetes Care* 30:234–238
 45. Heufelder AE, Saad F, Bunck MC, Gooren L 2009 Fifty-two-week treatment with diet and exercise plus transdermal testosterone reverses the metabolic syndrome and improves glycemic control in men with newly diagnosed type 2 diabetes and subnormal plasma testosterone. *J Androl* 30:726–733
 46. Gopal RA, Bothra N, Acharya SV, Ganesh HK, Bandgar TR, Menon PS, Shah NS 2010 Treatment of hypogonadism with testosterone in patients with type 2 diabetes mellitus. *Endocr Pract* 16:570–576
 47. Ding EL, Song Y, Malik VS, Liu S 2006 Sex differences of endogenous sex hormones and risk of type 2 diabetes: a systematic review and meta-analysis. *JAMA* 295:1288–1299
 48. Yialamas MA, Dwyer AA, Hanley E, Lee H, Pitteloud N, Hayes FJ 2007 Acute sex steroid withdrawal reduces insulin sensitivity in healthy men with idiopathic hypogonadotropic hypogonadism. *J Clin Endocrinol Metab* 92:4254–4259
 49. Rabiee A, Dwyer AA, Caronia LM, Hayes FJ, Yialamas MA, Andersen DK, Thomas B, Torriani M, Elahi D 2010 Impact of acute biochemical castration on insulin sensitivity in healthy adult men. *Biochem Res* 35:71–84
 50. Lapauw B, Ouwens M, 't Hart LM, Wuys B, Holst JJ, T'Sjoen G, Kaufman JM, Ruige JB 2010 Sex steroids affect triglyceride handling, glucose-dependent insulinotropic polypeptide, and insulin sensitivity: a 1-week randomized clinical trial in healthy young men. *Diabetes Care* 33:1831–1833
 51. Bojesen A, Birkebaek N, Kristensen K, Heickendorff L, Mosekilde L, Christiansen JS, Gravholt CH 2011 Bone mineral density in Klinefelter syndrome is reduced and primarily determined by muscle strength and resorptive markers, but not directly by testosterone. *Osteoporos Int* 22:1441–1450
 52. Maggio M, Ceda GP, Lauretani F, Bandinelli S, Metter EJ, Guralnik JM, Basaria S, Cattabiani C, Luci M, Dall'Aglio E, Vignali A, Volpi R, Valenti G, Ferrucci L 2011 Gonadal status and physical performance in older men. *Aging Male* 14:42–47
 53. Page ST, Amory JK, Bowman FD, Anawalt BD, Matsumoto AM, Bremner WJ, Tenover JL 2005 Exogenous testosterone (T) alone or with finasteride increases physical performance, grip strength, and lean body mass in older men with low serum T. *J Clin Endocrinol Metab* 90:1502–1510
 54. Sheffield-Moore M, Dillon EL, Casperson SL, Gilkison CR, Padon-Jones D, Durham WJ, Grady JJ, Urban RJ 2011 A randomized

- pilot study of monthly cycled testosterone replacement or continuous testosterone replacement versus placebo in older men. *J Clin Endocrinol Metab* 96:E1831–E1837
55. Srinivas-Shankar U, Roberts SA, Connolly MJ, O'Connell MD, Adams JE, Oldham JA, Wu FC 2010 Effects of testosterone on muscle strength, physical function, body composition, and quality of life in intermediate-frail and frail elderly men: a randomized, double-blind, placebo-controlled study. *J Clin Endocrinol Metab* 95:639–650
 56. Aksglaede L, Skakkebaek NE, Almstrup K, Juul A 2011 Clinical and biological parameters in 166 boys, adolescents and adults with nonmosaic Klinefelter syndrome: a Copenhagen experience. *Acta Paediatr* 100:793–806
 57. Smyth CM, Bremner WJ 1998 Klinefelter syndrome. *Arch Intern Med* 158:1309–1314
 58. Aksglaede L, Molgaard C, Skakkebaek NE, Juul A 2008 Normal bone mineral content but unfavourable muscle/fat ratio in Klinefelter syndrome. *Arch Dis Child* 93:30–34
 59. Singh R, Artaza JN, Taylor WE, Gonzalez-Cadavid NF, Bhasin S 2003 Androgens stimulate myogenic differentiation and inhibit adipogenesis in C3H10T1/2 pluripotent cells through an androgen receptor-mediated pathway. *Endocrinology* 144:5081–5088
 60. Tsai EC, Boyko EJ, Leonetti DL, Fujimoto WY 2000 Low serum testosterone level as a predictor of increased visceral fat in Japanese-American men. *Int J Obes Relat Metab Disord* 24:485–491
 61. Mårin P, Holmäng S, Jönsson L, Sjöström L, Kvist H, Holm G, Lindstedt G, Björntorp P 1992 The effects of testosterone treatment on body composition and metabolism in middle-aged obese men. *Int J Obes Relat Metab Disord* 16:991–997
 62. Pasquali R, Casimirri F, Cantobelli S, Melchionda N, Morselli Labate AM, Fabbri R, Capelli M, Bortoluzzi L 1991 Effect of obesity and body fat distribution on sex hormones and insulin in men. *Metabolism* 40:101–104
 63. Wang C, Jackson G, Jones TH, Matsumoto AM, Nehra A, Perelman MA, Swerdloff RS, Traish A, Zitzmann M, Cunningham G 2011 Low testosterone associated with obesity and the metabolic syndrome contributes to sexual dysfunction and cardiovascular disease risk in men with type 2 diabetes. *Diabetes Care* 34:1669–1675
 64. Luisetto G, Mastrogiacomo I, Bonanni G, Pozzan G, Botteon S, Tizian L, Galuppo P 1995 Bone mass and mineral metabolism in Klinefelter's syndrome. *Osteoporos Int* 5:455–461
 65. Behre HM, Kliesch S, Leifke E, Link TM, Nieschlag E 1997 Long-term effect of testosterone therapy on bone mineral density in hypogonadal men. *J Clin Endocrinol Metab* 82:2386–2390
 66. Stepan JJ, Burckhardt P, Hána V 2003 The effects of three-month intravenous ibandronate on bone mineral density and bone remodeling in Klinefelter's syndrome: the influence of vitamin D deficiency and hormonal status. *Bone* 33:589–596
 67. Smals AG, Kloppenborg PW, Lequin RL, Beex L, Ross A, Benraad TJ 1977 The pituitary-thyroid axis in Klinefelter's syndrome. *Acta Endocrinol (Copenh)* 84:72–79
 68. Bjørn AM, Bojesen A, Gravholt CH, Laurberg P 2009 Hypothyroidism secondary to hypothalamic-pituitary dysfunction may be part of the phenotype in Klinefelter syndrome: a case-control study. *J Clin Endocrinol Metab* 94:2478–2481
 69. Haka-Ikse K, Stewart DA, Cripps MH 1978 Early development of children with sex chromosome aberrations. *Pediatrics* 62:761–766
 70. Ratcliffe SG, Axworthy D, Ginsborg A 1979 The Edinburgh study of growth and development in children with sex chromosome abnormalities. *Birth Defects Orig Artic Ser* 15:243–260
 71. Tennes K, Puck M, Orfanakis D, Robinson A 1977 The early childhood development of 17 boys with sex chromosome anomalies: a prospective study. *Pediatrics* 59:574–583
 72. Walzer S, Graham Jr JM, Bashir AS, Silbert AR 1982 Preliminary observations on language and learning in XXY boys. *Birth Defects Orig Artic Ser* 18:185–192
 73. Leonard MF, Sparrow S 1986 Prospective study of development of children with sex chromosome anomalies: New Haven Study IV. *Adolescence. Birth Defects Orig Artic Ser* 22:221–249
 74. Ratcliffe S 1999 Long-term outcome in children of sex chromosome abnormalities. *Arch Dis Child* 80:192–195
 75. Walzer S 1985 X chromosome abnormalities and cognitive development: implications for understanding normal human development. *J Child Psychol Psychiatry* 26:177–184
 76. Bender BG, Linden MG, Robinson A 1993 Neuropsychological impairment in 42 adolescents with sex chromosome abnormalities. *Am J Med Genet* 48:169–173
 77. Bender BG, Harmon RJ, Linden MG, Robinson A 1995 Psychosocial adaptation of 39 adolescents with sex chromosome abnormalities. *Pediatrics* 96:302–308
 78. Graham Jr JM, Bashir AS, Stark RE, Silbert A, Walzer S 1988 Oral and written language abilities of XXY boys: implications for anticipatory guidance. *Pediatrics* 81:795–806
 79. Bender BG, Puck MH, Salbenblatt JA, Robinson A 1986 Dyslexia in 47,XXY boys identified at birth. *Behav Genet* 16:343–354
 80. van Rijn S, Aleman A, Swaab H, Krijn T, Vingerhoets G, Kahn R 2007 What it is said versus how it is said: comprehension of affective prosody in men with Klinefelter (47,XXY) syndrome. *J Int Neuropsychol Soc* 13:1065–1070
 81. Pennington BF, Bender B, Puck M, Salbenblatt J, Robinson A 1982 Learning disabilities in children with sex chromosome anomalies. *Child Dev* 53:1182–1192
 82. Linden MG, Bender BG 2002 Fifty-one prenatally diagnosed children and adolescents with sex chromosome abnormalities. *Am J Med Genet* 110:11–18
 83. Rovet J, Netley C, Keenan M, Bailey J, Stewart D 1996 The psychoeducational profile of boys with Klinefelter syndrome. *J Learn Disabil* 29:180–196
 84. Bender BG, Linden MG, Harmon RJ 2001 Life adaptation in 35 adults with sex chromosome abnormalities. *Genet Med* 3:187–191
 85. Barker TE, Black FW 1976 Klinefelter syndrome in a military population. *Electroencephalographic, endocrine, and psychiatric status. Arch Gen Psychiatry* 33:607–610
 86. Fales CL, Knowlton BJ, Holyoak KJ, Geschwind DH, Swerdloff RS, Gonzalo IG 2003 Working memory and relational reasoning in Klinefelter syndrome. *J Int Neuropsychol Soc* 9:839–846
 87. Geschwind DH, Gregg J, Boone K, Karrim J, Pawlikowska-Haddad A, Rao E, Ellison J, Ciccodicola A, D'Urso M, Woods R, Rappold GA, Swerdloff R, Nelson SF 1998 Klinefelter's syndrome as a model of anomalous cerebral laterality: testing gene dosage in the X chromosome pseudoautosomal region using a DNA microarray. *Dev Genet* 23:215–229
 88. Giedd JN, Clasen LS, Wallace GL, Lenroot RK, Lerch JP, Wells EM, Blumenthal JD, Nelson JE, Tossell JW, Stayer C, Evans AC, Samango-Sprouse CA 2007 XXY (Klinefelter Syndrome): a pediatric quantitative brain magnetic resonance imaging case-control study. *Pediatrics* 119:e232–e240
 89. Itti E, Gaw Gonzalo IT, Pawlikowska-Haddad A, Boone KB, Mlikotic A, Itti L, Mishkin FS, Swerdloff RS 2006 The structural brain correlates of cognitive deficits in adults with Klinefelter's syndrome. *J Clin Endocrinol Metab* 91:1423–1427
 90. Kompus K, Westerhausen R, Nilsson LG, Hugdahl K, Jongstra S, Berglund A, Arver S, Savic I 2011 Deficits in inhibitory executive functions in Klinefelter (47, XXY) syndrome. *Psychiatry Res* 189:135–140
 91. Lee NR, Wallace GL, Clasen LS, Lenroot RK, Blumenthal JD, White SL, Celano MJ, Giedd JN 2011 Executive function in young males with Klinefelter (XXY) syndrome with and without comorbid attention-deficit/hyperactivity disorder. *J Int Neuropsychol Soc* 22:1–9
 92. Tartaglia NR, Ayari N, Hutaff-Lee C, Boada R 2012 Attention-deficit hyperactivity disorder symptoms in children and adolescents with sex chromosome aneuploidy: XXY, XXX, XYY, and XYYY. *J Dev Behav Pediatr* 33:309–318
 93. van Rijn S, Aleman A, Swaab H, Kahn R 2006 Klinefelter's syn-

- drome (karyotype 47,XXY) and schizophrenia-spectrum pathology. *Br J Psychiatry* 189:459–460
94. van Rijn S, Swaab H, Aleman A, Kahn RS 2008 Social behavior and autism traits in a sex chromosomal disorder: Klinefelter (47XXY) syndrome. *J Autism Dev Disord* 38:1634–1641
 95. DeLisi LE, Maurizio AM, Svetina C, Ardekani B, Szulc K, Nierenberg J, Leonard J, Harvey PD 2005 Klinefelter's syndrome (XXY) as a genetic model for psychotic disorders. *Am J Med Genet B Neuropsychiatr Genet* 135B:15–23
 96. Temple CM, Sanfilippo PM 2003 Executive skills in Klinefelter's syndrome. *Neuropsychologia* 41:1547–1559
 97. Boone KB, Swerdloff RS, Miller BL, Geschwind DH, Razani J, Lee A, Gonzalo IG, Haddad A, Rankin K, Lu P, Paul L 2001 Neuropsychological profiles of adults with Klinefelter syndrome. *J Int Neuropsychol Soc* 7:446–456
 98. Geschwind DH, Boone KB, Miller BL, Swerdloff RS 2000 Neurobehavioral phenotype of Klinefelter syndrome. *Ment Retard Dev Disabil Res Rev* 6:107–116
 99. Bender BG, Linden MG, Harmon RJ 2001 Neuropsychological and functional cognitive skills of 35 unselected adults with sex chromosome abnormalities. *Am J Med Genet* 102:309–313
 100. Nielsen J, Pelsen B 1987 Follow-up 20 years later of 34 Klinefelter males with karyotype 47,XXY and 16 hypogonadal males with karyotype 46,XY. *Hum Genet* 77:188–192
 101. Anell AL, Gustavson KH, Tenstam J 1970 Symptomatology in schoolboys with positive sex chromatin (the Klinefelter syndrome). *Acta Psychiatr Scand* 46:71–80
 102. Patwardhan AJ, Eliez S, Bender B, Linden MG, Reiss AL 2000 Brain morphology in Klinefelter syndrome: extra X chromosome and testosterone supplementation. *Neurology* 54:2218–2223
 103. Ross JL, Roeltgen DP, Stefanatos G, Benecke R, Zeger MP, Kushner H, Ramos P, Elder FF, Zinn AR 2008 Cognitive and motor development during childhood in boys with Klinefelter syndrome. *Am J Med Genet A* 146A:708–719
 104. Stemkens D, Roza T, Verrij L, Swaab H, van Werkhoven MK, Alizadeh BZ, Sinke RJ, Giltay JC 2006 Is there an influence of X-chromosomal imprinting on the phenotype in Klinefelter syndrome? A clinical and molecular genetic study of 61 cases. *Clin Genet* 70:43–48
 105. Vawter MP, Harvey PD, DeLisi LE 2007 Dysregulation of X-linked gene expression in Klinefelter's syndrome and association with verbal cognition. *Am J Med Genet B Neuropsychiatr Genet* 144B:728–734
 106. DeLisi LE, Friedrich U, Wahlstrom J, Boccio-Smith A, Forsman A, Eklund K, Crow TJ 1994 Schizophrenia and sex chromosome anomalies. *Schizophr Bull* 20:495–505
 107. Boks MP, de Vette MH, Sommer IE, van Rijn S, Giltay JC, Swaab H, Kahn RS 2007 Psychiatric morbidity and X-chromosomal origin in a Klinefelter sample. *Schizophr Res* 93:399–402
 108. Bruining H, Swaab H, Kas M, van Engeland H 2009 Psychiatric characteristics in a self-selected sample of boys with Klinefelter syndrome. *Pediatrics* 123:e865–e870
 109. Bishop DV, Jacobs PA, Lachlan K, Wellesley D, Barnicoat A, Boyd PA, Fryer A, Middlemiss P, Smithson S, Metcalfe K, Shears D, Leggett V, Nation K, Scerif G 2011 Autism, language and communication in children with sex chromosome trisomies. *Arch Dis Child* 96:954–959
 110. Ratcliffe SG, Butler GE, Jones M 1990 Edinburgh study of growth and development of children with sex chromosome abnormalities. IV. *Birth Defects Orig Artic Ser* 26:1–44
 111. Nielsen J, Pelsen B, Sørensen K 1988 Follow-up of 30 Klinefelter males treated with testosterone. *Clin Genet* 33:262–269
 112. Simm PJ, Zacharin MR 2006 The psychosocial impact of Klinefelter syndrome—a 10 year review. *J Pediatr Endocrinol Metab* 19:499–505
 113. Wang C, Swerdloff RS, Iranmanesh A, Dobs A, Snyder PJ, Cunningham G, Matsumoto AM, Weber T, Berman N 2000 Transdermal testosterone gel improves sexual function, mood, muscle strength, and body composition parameters in hypogonadal men. Testosterone Gel Study Group. *J Clin Endocrinol Metab* 85:2839–2853
 114. Cherrier MM, Asthana S, Plymate S, Baker L, Matsumoto AM, Peskind E, Raskind MA, Brodtkin K, Bremner W, Petrova A, La-Tendresse S, Craft S 2001 Testosterone supplementation improves spatial and verbal memory in healthy older men. *Neurology* 57:80–88
 115. Bhasin S, Cunningham GR, Hayes FJ, Matsumoto AM, Snyder PJ, Swerdloff RS, Montori VM 2006 Testosterone therapy in adult men with androgen deficiency syndromes: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 91:1995–2010
 116. Aksglaede L, Andersson AM, Jørgensen N, Jensen TK, Carlsen E, McLachlan RI, Skakkebaek NE, Petersen JH, Juul A 2007 Primary testicular failure in Klinefelter's syndrome: the use of bivariate luteinizing hormone-testosterone reference charts. *Clin Endocrinol (Oxf)* 66:276–281
 117. Srinivas-Shankar U, Wu FC 2006 Drug insight: testosterone preparations. *Nat Clin Pract Urol* 3:653–665
 118. Rogol AD, Tartaglia N 2010 Considerations for androgen therapy in children and adolescents with Klinefelter syndrome (47, XXY). *Pediatr Endocrinol Rev* 8(Suppl 1):145–150
 119. Lue Y, Liu PY, Erkkila K, Ma K, Schwarcz M, Wang C, Swerdloff RS 2010 Transplanted XY germ cells produce spermatozoa in testes of XXY mice. *Int J Androl* 33:581–587
 120. Liu PY, Kalak R, Lue Y, Jia Y, Erkkila K, Zhou H, Seibel MJ, Wang C, Swerdloff RS, Dunstan CR 2010 Genetic and hormonal control of bone volume, architecture, and remodeling in XXY mice. *J Bone Miner Res* 25:2148–2154
 121. Lewejohann L, Damm OS, Luetjens CM, Hämäläinen T, Simoni M, Nieschlag E, Gromoll J, Wistuba J 2009 Impaired recognition memory in male mice with a supernumerary X chromosome. *Physiol Behav* 96:23–29
 122. Hollegaard MV, Grauholm J, Børglum A, Nyegaard M, Nørgaard-Pedersen B, Ørntoft T, Mortensen PB, Wiuf C, Mors O, Didriksen M, Thorsen P, Hougaard DM 2009 Genome-wide scans using archived neonatal dried blood spot samples. *BMC Genomics* 10:297
 123. Vorona E, Zitzmann M, Gromoll J, Schüring AN, Nieschlag E 2007 Clinical, endocrinological, and epigenetic features of the 46,XX male syndrome, compared with 47,XXY Klinefelter patients. *J Clin Endocrinol Metab* 92:3458–3465
 124. van den Bergh JP, Hermus AR, Spruyt AI, Sweep CG, Corstens FH, Smals AG 2001 Bone mineral density and quantitative ultrasound parameters in patients with Klinefelter's syndrome after long-term testosterone substitution. *Osteoporos Int* 12:55–62
 125. Stewart DA, Netley CT, Park E 1982 Summary of clinical findings of children with 47,XXY, 47,XYY, and 47,XXX karyotypes. *Birth Defects Orig Artic Ser* 18:1–5
 126. Fricke GR, Mattern HJ, Schweikert HU, Schwanitz G 1984 Klinefelter's syndrome and mitral valve prolapse. An echocardiographic study in twenty-two patients. *Biomed Pharmacother* 38:88–97
 127. Andersen NH, Bojesen A, Kristensen K, Birkebaek NH, Fedder J, Bennet P, Christiansen JS, Gravholt CH 2008 Left ventricular dysfunction in Klinefelter syndrome is associated to insulin resistance, abdominal adiposity and hypogonadism. *Clin Endocrinol (Oxf)* 69:785–791
 128. Hultborn R, Hanson C, Köpf I, Verbiené I, Warnhammar E, Weimarck A 1997 Prevalence of Klinefelter's syndrome in male breast cancer patients. *Anticancer Res* 17:4293–4297